

Lattice QCD signal for a bottom-bottom tetraquark

Pedro Bicudo*

Dep. Física and CFTP, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

Marc Wagner†

*Johann Wolfgang Goethe-Universität Frankfurt am Main, Institut für Theoretische Physik,
Max-von-Laue-Straße 1, D-60438 Frankfurt am Main, Germany and
European Twisted Mass Collaboration (ETMC)*

Utilizing lattice QCD results for the potential of two static antiquarks and two dynamical quarks as well as quark model techniques for the dynamics of two heavy antiquarks in a cloud of two light quarks, we are provided with an accurate framework for the study of possibly existing heavy-heavy-light-light tetraquarks. Among the possible quantum numbers of such a system, we find binding in only one channel, the scalar isosinglet. Solving the Schrödinger equation for the displacement of the heavy antiquarks and taking systematic errors into account, we find an antibottom-antibottom-light-light bound state with a confidence level of around $1.8\sigma \dots 3.0\sigma$ and binding energy of approximately $30 \text{ MeV} \dots 57 \text{ MeV}$.

PACS numbers: 12.38.Gc, 13.75.Lb, 14.40.Rt, 14.65.Fy.

Experimentally exotic hadrons have been searched for many years, because as soon as quarks were proposed in the sixties, it became clear that systems more complex than mesons and baryons could possibly exist. However, exotic hadrons are very elusive systems. They are much harder to observe experimentally, to understand by theoretical model calculations, and to simulate by means of lattice QCD than the conventional mesons and baryons. To confirm the existence or non-existence of exotic hadrons still remains an important problem in QCD.

A frequently discussed exotic multiquark is the tetraquark. It was already proposed in the seventies [1] as a bound state formed by two quarks and two antiquarks. There are several hadronic resonances, which are tetraquark candidates, e.g. σ , κ , D_{s0}^* or D_{s1} [2]. The most recent tetraquark candidate has been claimed by the Belle Collaboration [3], observing in five different $\Upsilon(5S)$ decay channels two new charged bottomonium resonances Z_b with masses 10610 MeV and 10650 MeV and narrow widths of the order of 15 MeV , where the charge can only come from the presence of a light quark and a light antiquark. However, the tetraquark nature of these resonance is disputed [4].

Notice that the experimental particle physics collaborations are technically improving rather impressively. One decade ago SELEX at FNAL already studied the doubly charmed baryons. Presently, BELLE at KEK, CDF and DØ at FNAL, and LHCb at CERN have already observed bottom hadrons. Thus, they may possibly be able to search for charm-bottom and bottom-bottom tetraquarks, if a strong evidence is presented by theoretical calculations.

Tetraquark studies face several difficulties. (1) Mesons

and baryons only decay strongly, when the confining string breaks, a quark antiquark pair is created and either two mesons or a meson and a baryon are formed. In contrast to that, tetraquarks are directly open to meson-meson decay. (2) Moreover, tetraquarks are relativistic four-body systems, which are highly complex few-body systems. (3) And on the top of these technical difficulties no model (since the onset of QCD [1] up to the present) seems to be sufficiently well calibrated to address multiquark binding: different quark models, even when producing similar meson or baryon spectra, usually differ significantly in predictions for tetraquarks.

A strategy to avoid many technical difficulties associated with tetraquark studies consists in searching for bound states rather than for resonances, which is e.g. appropriate, when two heavy antiquarks (or equivalently two heavy quarks) are involved. This strategy was already identified in the eighties [5]. On the one hand, it is plausible that any light tetraquark can only be treated as a resonance, because it couples to two-meson channels with identical quantum numbers, where at least one of the mesons is a pion or kaon. Due to chiral symmetry, these are very light mesons and, consequently, the tetraquark will have a rather light open decay channel. On the other hand the presence of two heavy antiquarks is sufficient to force all involved hadrons, i.e. the tetraquark and any two-meson channels with the same quantum numbers, to escape chiral symmetry. Moreover, the heavy antiquarks also reduce the technical complexity of the four quark system, since heavy antiquarks are non-relativistic. This in turn allows for the Born-Oppenheimer approximation: for the light quarks the heavy antiquarks can be regarded as static color charges; once the light diquark energy is determined, it can be utilized as an effective potential for the heavy antiquarks.

The only remaining theoretical problem is to obtain the correct effective potential, which has been studied by lattice QCD methods mainly in the quenched approxima-

*Electronic address: bicudo@ist.utl.pt

†Electronic address: mwagner@th.physik.uni-frankfurt.de

tion (cf. e.g. [6, 7] and references therein). Only recently computations of this potential with dynamical sea quarks have been performed [8–10]. Here we use such dynamical results obtained with a comprehensive set of four quark operators [8, 10]. While for the static antiquarks the only relevant variable is their separation, the two light u/d quarks can be combined in $8 \times 8 = 64$ different ways (each light quark has two isospin, two spin and two parity degrees of freedom). Symmetries and quantum numbers of BB states are explained in detail in [8]. In this work we focus on the two attractive channels between ground state static-light mesons (B mesons): a scalar isosinglet (cf. Fig. 1(a)), which is most attractive, and a vector isotriplet (cf. Fig. 1(b)), which is less attractive.

To motivate an ansatz to fit the lattice results for these potentials, first remember that the pair of heavy antiquarks is immersed in a cloud of two light quarks. The size of this cloud is crucial for the heavy antiquark-antiquark interaction. When the two antiquarks are much closer than twice the typical light quark cloud radius in a heavy-light meson, the antiquark-antiquark interaction is a typical diquark interaction (cf. Fig. 2(a)). However, when the separation of the two antiquarks becomes larger, the light quarks screen the heavy antiquark charges as in Fig. 2(b) and the antiquark-antiquark interaction vanishes.




Figure 1: (Color online). The static antiquark-antiquark potential as a function of the separation in units of the lattice spacing $a \approx 0.079$ fm. (a) Three parameter fit (α , d , p) of ansatz (1) to the most attractive channel, the scalar isosinglet. (b) Two parameter fit (α , d ; $p = 2.0$ fixed) of ansatz (1) to the less attractive vector isotriplet.




Figure 2: (Color online). Screening of the antiquark flux tube interaction. We show two scenarios: (a) the heavy antiquarks are close and the screening of the light quarks has little effect on the antiquark-antiquark interaction; (b) the heavy quarks are well separated and their color charge is totally screened by the light quark wavefunctions.

The diquark interaction for systems containing only static quarks has also been studied by lattice QCD. Flux tubes have been observed for static baryons, static tetraquarks and static pentaquarks [11, 12]. It seems established that at large separations r the diquark potential is linear and confining $\sim \sigma r$. At small separations r the potential is rather Coulomb-like, i.e. $\sim -\alpha/r$. A similar system, which has been studied in even more detail by lattice methods is the “static-static meson” or static potential; here $\alpha \approx 0.3 \dots 0.4$, while $\sigma \approx (0.44 \text{ GeV})^2$ is estimated from quark model fits and often used to set the scale. For two heavy antiquarks in a cloud of two light quarks we expect a similar Coulomb-like potential of order $-\alpha/r$ at small separations. At larger separations (in contrast to the purely static case) the potential should be screened by the light quarks, as discussed in the previous paragraph and illustrated in Fig. 2(b).

The screening of the heavy color charge is due to the decrease of the wave function ψ of the light quark with respect to its separation from the heavy quark. One expects this decrease to follow an exponential of a power law, i.e. $\psi(r) \propto \exp(-(r/d)^p)$, where d characterizes the size of the quark-antiquark system, i.e. a B meson. If the quark-antiquark interaction inside a B meson is dominated by a Coulomb-like term, the wave function is similar to that of a hydrogen atom, i.e. $p = 1$. In case the potential is rather linear, the non-relativistic Schrödinger equation is solved by Airy functions corresponding to $p = 3/2$. A similar, but relativistic treatment of the light quark yields $p = 2$ instead.

The above considerations suggest the following ansatz to model the heavy antiquark-antiquark potential:

$$V(r) = -\frac{\alpha}{r} \exp \left[-\left(\frac{r}{d} \right)^p \right], \quad (1)$$

where it is expected that $\alpha \approx 0.3 \dots 0.4$, d is around half the size of a B meson, i.e. $d \lesssim 0.5$ fm and $p \approx 1.0 \dots 2.0$.

We perform χ^2 minimizing fits of the ansatz (1) to the lattice results for the heavy antiquark-antiquark interactions shown in Fig. 1. Notice that data points for separations $r/a = 0$ and $r/a = 1$ are excluded from the fits, because they suffer from lattice discretization errors. For $r/a \geq 2$ it has been checked, that these discretization errors are negligible compared to the statistical errors by

using different static quark actions [8]. On the other hand, data points for large separations have little relevance since the potential vanishes exponentially fast, due to screening, while the statistical error remains similar. Thus we utilize the points at distances $r/a = 2, 3, 4, 5, 6$ for our fits.

For the scalar isosinglet we are able to determine all three parameters α , d and p via fitting. For the vector isotriplet a three parameter fit is not stable; therefore, we only fit two parameters, α and d , while fixing the exponent to its expected value $p = 2.0$. The fits are also shown in Fig. 1, while numerical results are collected in Table I (statistical errors for α , d and p have been determined via an elaborate Jackknife analysis starting on the level of the bare lattice correlation functions). The uncorrelated $\chi^2/\text{dof} \ll 1$ indicates consistency of the lattice data and our ansatz (1). Moreover, note that the resulting values for α , d and p are in agreement with phenomenological expectations.

Table I: χ^2 minimizing fit results of the ansatz (1) to the lattice static antiquark-antiquark potential; fitting range $2 \leq r/a \leq 6$; lattice spacing $a \approx 0.079$ fm

channel	α	d/a	p	χ^2/dof
scalar isosinglet	0.293(33)	4.51(54)	2.74(1.20)	0.35
vector isotriplet	0.201(77)	2.48(69)	2.0 (fixed)	0.06

The potential of Eq. (1) with the fit parameters from Table I corresponds to the energy of a static-static-light-light four quark system minus the energy of a pair of static-light ground state mesons. To obtain the energy of a heavy-heavy-light-light system, where the antiquarks have a heavy, but finite mass, one also needs to consider a kinetic term for the heavy antiquarks resulting in the Hamiltonian

$$H = \frac{\mathbf{p}^2}{2\mu} + 2m_B + V(r), \quad (2)$$

where μ is the reduced antiquark mass.

Notice that, because of screening, at large separations each heavy antiquark carries the mass of a B meson and thus $\mu = m_B/2$, whereas at small separations it carries just the energy of a heavy quark $\mu = m_b/2$. When investigating the existence of four quark bound states, we consider both mass values, which differ by around 5% ($m_B = 5279$ MeV [2], $m_b = 4977$ MeV in quark models [13]). Another possible source of systematic error is associated with the physical value of the lattice spacing. This error is introduced, when converting the “size parameter” d from dimensionless lattice units to physical units. We investigate the magnitude of this error, by using values for the lattice spacing determined by rather different scale setting procedures: $a = 0.079$ fm is used in many ETMC publications and is obtained from m_π , f_π and chiral perturbation theory [14], while $a = 0.096$ fm corresponds to identifying the lattice result for the Sommer parameter r_0 with its typical value 0.5 fm.

To get an analytical qualitative understanding, we first derive an approximate rule for the existence/non-existence of a bound state applying the Bohr-Sommerfeld relation corrected by the WKB approximation, to include the zero point energy of $1/2$. For the radial equation in three dimensions we get,

$$4 \int_{c_1}^{c_2} dr \sqrt{2\mu \left(E - V(r) - \frac{l(l+1)}{2\mu r^2} \right)} = 2\pi \left(n + \frac{3}{2} \right), \quad (3)$$

where c_1 and c_2 are the classical turning points. Inserting (1), specializing to angular momentum $l = 0$ and integrating, yields the condition for having at least one bound state,

$$\mu\alpha d \geq \frac{9\pi^2}{128 \times 2^{1/p} \Gamma^2(1 + 1/2p)}. \quad (4)$$

The right hand side of Eq. (4) has a rather moderate dependence on the exponent p . For example, when p increases from the expected values of 1.0 to 2.0, the right hand side only changes from 0.44 to 0.60. Thus the existence of a bound state mainly depends on the product of parameters $\mu\alpha d$. With the fit parameters for the scalar isosinglet from Table I and a bottom quark one roughly obtains $\mu\alpha d \approx 2.5 \text{ GeV} \times 0.29 \times 4.5 \times 0.079 \text{ fm} \approx 1.3 \gg 0.44 \dots 0.60$, which is a strong indication for the existence of a bound state. A similar calculation for the vector isotriplet yields $\mu\alpha d \approx 0.5$, i.e. the existence of a bound state in this channel is rather questionable.

Note, however, that these are only crude analytical estimates, because e.g. the WKB approximation is questionable, when the potential is divergent at the origin as for a Coulomb-like potential. To investigate the existence of a bound state rigorously, we numerically solve the Schrödinger equation with the Hamiltonian (2). The strongest binding is expected in an s-wave, for which the radial equation is

$$\left[-\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} + 2m_B + V(r) \right] R(r) = ER(r) \quad (5)$$

with the wave function $\psi = \psi(r) = R(r)/r$. We impose Dirichlet boundary conditions $R(r_{\text{max}}) = 0$ at sufficiently large r_{max} (we checked that results are stable for $r_{\text{max}} \gtrsim 10$ fm). The radial equation (5) can be solved by standard methods (e.g. 4th order Runge-Kutta shooting) up to arbitrary numerical precision.

In accordance with our analytical estimates we find binding for heavy bottom-bottom tetraquarks in the scalar isosinglet channel. The binding energy $E - 2m_B \approx 30 \text{ MeV} \dots 57 \text{ MeV}$ depends to some extent on the reduced mass (either $\mu = m_b/2$ or $\mu = m_B/2$) and on the value of the lattice spacing (either $a = 0.079$ fm or $a = 0.096$ fm); cf. Table II. Note that the binding energy is negative by around $1.76\sigma \dots 3.00\sigma$. This confirms the existence of an heavy-heavy-light-light tetraquark with a confidence level of $\gtrsim 96\%$. The probability to find the two heavy antiquarks at separation r is proportional to

Table II: Binding energy and confidence for the existence of a heavy-heavy-light-light tetraquark for the scalar isosinglet channel.

μ	a	$E - 2m_B$	confidence for binding
$m_b/2$	0.079 fm	-30(17) MeV	1.76σ , 96% binding
$m_b/2$	0.096 fm	-49(17) MeV	2.88σ , 100% binding
$m_B/2$	0.079 fm	-38(18) MeV	2.11σ , 98% binding
$m_B/2$	0.096 fm	-57(19) MeV	3.00σ , 100% binding




Figure 3: (Color online). Radial probability density $\propto |R(r)|^2$ for the separation of the heavy antiquarks (scalar isosinglet).

$|R(r)|^2$ and shown in Fig. 3. The average separation is around 0.25 fm.

Solving Schrödinger's equation (5) for the vector isotriplet gives strong indication that the potential of this channel is too weak to generate a bound state, i.e. a tetraquark. The resulting wave function is essentially a plane wave with positive energy $E - 2m_B$ within more than 10σ .

Further possible systematic errors are not expected to weaken the binding. The static approximation of a ground state with bottom antiquarks is valid, since the bottom quark mass is a very hard scale compared with the scale Λ_{QCD} . Moreover, lattice computations of B mesons [15, 16] and b baryons [17] within the same lattice setup showed that static quarks are a rather good approximation of bottom quarks. Finite volume effects are exponentially suppressed in lattice results and the spatial extension of the lattice $L \approx 1.9$ fm seems large compared to the typical size of the tetraquark, which is related to $d \approx 4.51 a \approx 0.36$ fm and the average separation of the heavy antiquarks, which is ≈ 0.25 fm. Finally, the light u/d quark masses in the lattice computation are unphysically heavy (corresponding to $m_\pi \approx 340$ MeV). Decreasing the light quark masses to their physical values should increase the light cloud radius of a heavy-light meson and, therefore, lead to stronger binding (cf. Fig. 2(a)).

To summarize, we find clearest evidence for the existence of an antibottom-antibottom-light-light tetraquark bound state. This result is very promising with respect to further lattice QCD calculations with higher statistics or even lighter dynamical quarks, and for experimental searches of doubly bottom hadrons.

Acknowledgments

P.B. thanks the hospitality of IFT and FCT grants CERN/FP/116383/2010, CERN/FP/123612/2011.

M.W. acknowledges support by the Emmy Noether Programme of the DFG (German Research Foundation), grant WA 3000/1-1.

-
- [1] R. L. Jaffe, Phys. Rev. D **15**, 267 (1977).
 - [2] J. Beringer et al. [Particle Data Group], Phys. Rev. D **86**, 010001 (2012).
 - [3] A. Bondar *et al.* [Belle Collaboration], Phys. Rev. Lett. **108**, 122001 (2012) [arXiv:1110.2251 [hep-ex]].
 - [4] D. V. Bugg, Europhys. Lett. **96**, 11002 (2011) [arXiv:1105.5492 [hep-ph]].
 - [5] J. L. Ballot and J. M. Richard, Phys. Lett. B **123**, 449 (1983).
 - [6] C. Michael and P. Penanen [UKQCD Collaboration], Phys. Rev. D **60**, 054012 (1999) [arXiv:hep-lat/9901007].
 - [7] W. Detmold, K. Orginos and M. J. Savage, Phys. Rev. D **76**, 114503 (2007) [arXiv:hep-lat/0703009].
 - [8] M. Wagner [ETM Collaboration], PoS LATTICE **2010**, 162 (2010) [arXiv:1008.1538 [hep-lat]].
 - [9] G. Bali and M. Hetzenegger, PoS LATTICE2010, 142 (2010) [arXiv:1011.0571 [hep-lat]].
 - [10] M. Wagner [ETM Collaboration], Acta Phys. Polon. Supp. **4**, 747 (2011) [arXiv:1103.5147 [hep-lat]].
 - [11] N. Cardoso, M. Cardoso and P. Bicudo, Phys. Rev. D **84**, 054508 (2011) [arXiv:1107.1355 [hep-lat]].
 - [12] N. Cardoso and P. Bicudo, arXiv:1209.1532 [hep-lat].
 - [13] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
 - [14] R. Baron *et al.* [ETM Collaboration], JHEP **1008**, 097 (2010) [arXiv:0911.5061 [hep-lat]].
 - [15] K. Jansen *et al.* [ETM Collaboration], JHEP **0812**, 058 (2008) [arXiv:0810.1843 [hep-lat]].
 - [16] C. Michael *et al.* [ETM Collaboration], JHEP **1008**, 009 (2010) [arXiv:1004.4235 [hep-lat]].
 - [17] M. Wagner *et al.* [ETM Collaboration], JHEP **1107**, 016 (2011) [arXiv:1104.4921 [hep-lat]].